

*Nativity and Cardiovascular  
Dysregulation: Evidence from the 2001–  
2016 National Health and Nutrition  
Examination Survey*

**Lauren A. Doamekpor, Jessica  
L. Gleason, Ijeoma Opara & Ndidiamaka  
N. Amutah-Onukagha**

**Journal of Racial and Ethnic Health  
Disparities**

ISSN 2197-3792

J. Racial and Ethnic Health Disparities  
DOI 10.1007/s40615-020-00765-z



**Your article is protected by copyright and all rights are held exclusively by W. Montague Cobb-NMA Health Institute. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".**



# Nativity and Cardiovascular Dysregulation: Evidence from the 2001–2016 National Health and Nutrition Examination Survey

Lauren A. Doamekpor<sup>1</sup> · Jessica L. Gleason<sup>2</sup> · Ijeoma Opara<sup>3</sup> · Ndidiamaka N. Amutah-Onukagha<sup>4</sup>

Received: 11 November 2019 / Revised: 14 March 2020 / Accepted: 22 April 2020  
© W. Montague Cobb-NMA Health Institute 2020

## Abstract

**Objectives** To examine nativity-based differences in 3 cardiovascular biomarkers commonly used to assess cardiovascular dysregulation.

**Methods** Data was pooled from the 2001–2016 National Health and Nutrition Examination Survey to compare biomarker risk scores for the US-born ( $n = 4693$ ) and foreign-born ( $n = 2968$ ) Black adults. We used multivariable-adjusted logistic regression to assess the association between nativity and cardiovascular biomarkers, controlling for gender, age, health behaviors, and socio-economic status.

**Results** In the full model, a foreign-born health advantage was not observed in all 3 cardiovascular biomarkers. In fact, foreign-born Blacks were almost twice as likely to have high mean diastolic blood pressure compared with the US-born individuals (OR = 1.82; 95% CI = 1.15, 2.88) and had an increased risk of high 60-s pulse. Foreign-born individuals living in the USA for less than 5 years were 62% less likely to have high mean systolic blood pressure than individuals living in the USA for 20 years or more.

**Conclusions** The foreign-born health advantage among Blacks was not observed in the cardiovascular biomarkers under study, suggesting that the commonly cited Healthy Immigrant Effect may need to be reassessed.

**Keywords** Nativity · Cardiovascular risk · Biomarkers

✉ Lauren A. Doamekpor  
doamekpor3@gmail.com

Jessica L. Gleason  
jgleason@terpmail.umd.edu

Ijeoma Opara  
ijeoma.opara@stonybrook.edu

Ndidiamaka N. Amutah-Onukagha  
ndidiamaka.amutah\_onukagha@tufts.edu

<sup>1</sup> Guidehouse LLP, Washington, DC 20006, USA

<sup>2</sup> Department of Family Science, School of Public Health, University of Maryland, College Park, MD, USA

<sup>3</sup> Stony Brook University School of Social Welfare, Stony Brook, NY 11794, USA

<sup>4</sup> Department of Public Health and Community Medicine, School of Medicine, Tufts University, 136 Harrison Avenue, Boston, MA 02111, USA

## Introduction

The Healthy Immigrant Effect (HIE) describes the phenomenon in which people who migrate to the United States (US) experience better health outcomes than those born in the USA. Immigrants' cardiovascular risk profile has garnered special attention as cardiovascular disease rates increase in the USA and globally. Recently, attention has shifted to the health of Black immigrants as this population has seen rapid growth, increasing by over 800,000, an 11-fold increase, between 1970 and 2000. In 2015, 2.1 million Africans lived in the USA and this rapid growth is expected to increase [1, 2].

Cardiovascular health disparities between all non-Hispanic Blacks and Whites in the USA have been well-documented with the general consensus that non-Hispanic Blacks bear a disproportionately higher burden of cardiovascular disease than Whites [3, 4]. The risk of cardiovascular disease increases as blood pressure increases and puts individuals at risk for developing hypertension and other chronic diseases that lead to negative health outcomes. Hypertension, or chronically elevated blood pressure, is a major health concern for non-

Hispanic Blacks in particular, as this group has the highest rate of hypertension in the world [5]. Studies of nativity and hypertension indicate that the US-born Blacks have a higher risk of being diagnosed with hypertension and tend to be diagnosed at younger ages compared with non-Hispanic Whites and other ethnic minority groups [5]. However, an important limitation of these studies is the lack of an examination of nativity-based differences that consider the heterogeneity within the US Black population. An examination of these differences can provide valuable evidence to better understand disparities in this area.

Past views on the cardiovascular health of foreign-born Blacks were that only after living in the USA did they acquire high blood pressure, hypertension, and cardiovascular disease [6]. However, the prevalence of cardiovascular disease is on the rise on the African continent and other Black-majority nations [7–9], leading to a significant number of immigrants arriving to the USA with already compromised cardiovascular systems and disease. In many African countries, cardiovascular disease is the leading cause of death [6, 7], and the prevalence of risk factors for cardiovascular disease such as the diagnoses of diabetes and hypertension is also increasing in many African developing areas [10]. Hypertension is seen as the most common underlying risk factor for cardiovascular disease in Africa [11]. In 2025, adults living with hypertension are expected to increase by 60%, equating to approximately 1.56 billion total individuals that will be burdened by the disease [12].

Elevated levels of diastolic and systolic blood pressure (DBP) (SBP) are both indicators of hypertension and predictors of cardiovascular disease [13]. Emerging evidence has shown that the variability of SBP has the potential to affect estimation of an individual's 10-year cardiovascular disease risk with increases in SBP nearly doubling the overall risk of stroke and coronary heart disease mortality [14, 15]. Of these two indicators, SBP is often considered the more clinically relevant marker as a strong predictor of cardiovascular disease among hypertensives. However, DBP is a stronger predictor of cardiovascular morbidity among individuals under the age of 50 [16]. Thus, DBP may be an equally important and often overlooked indicator of differences in cardiovascular morbidity risk between foreign-born Blacks and the US-born Blacks although few studies have explored this association [17]. One study found that SBP levels are higher in the US-born Blacks compared with foreign-born Blacks from sub-Saharan Africa, but to our knowledge, there are no studies exploring nativity differences in DBP [18].

In comparing foreign-born with the US-born Blacks, O'Connor et al., found that foreign-born Blacks had higher fasting glucose, a higher prevalence of hypertension, and greater visceral adiposity, all of which increased their cardiometabolic risk [6]. Other studies have found that African immigrants had higher rates of hypertension than African

Americans [19–21]. Alternatively, a recent study found that foreign-born Blacks were close to 40% less likely to have hypertension than their US-born counterparts [17].

Studies examining the HIE provide insight into potential protective factors from the risk of cardiovascular disease. Foreign-born Blacks are more likely to have a higher SES and higher education level upon arrival, thus providing them with access to quality healthcare that might help to mitigate health risks [22]. When the foreign-born Black group is disaggregated, research shows that Caribbean-born Blacks have significantly lower rates of cardiovascular death and better health outcomes than the US-born Blacks [22]. However, when compared with West African-born Blacks, Caribbean-born Blacks had higher average systolic and diastolic blood pressures and had a higher prevalence of hypertension [23].

Given these inconsistent findings, it is important to understand how nativity might impact cardiovascular risk by examining the individual components of hypertension; SBP, DBP, and 60-s pulse. While hypertension is a composite measure of SBP and DBP, our study examines these components separately to determine what may drive differences in these biomarkers, and how sociodemographic variables influence associations between nativity and individual biomarkers. Several large epidemiological studies, such as the Framingham Heart Study, have suggested that SBP and pulse may have more influence on blood pressure than DBP [24]. To our knowledge, no other study has examined nativity-based differences in individual cardiovascular biomarkers. The objective of the current study is to examine the nativity-based differences in cardiovascular biomarkers and assess the sociodemographic variables that may influence these factors. We also examine how the sociodemographic variables under study work differently in the US-born and foreign-born populations.

## Methods

We pooled data from the 2001–2016 National Health and Nutrition Examination Survey (NHANES) to create the analytical sample. NHANES uses a stratified, multi-stage probability sample to provide national estimates of the nutritional status and health for the civilian, noninstitutionalized population of the USA [25]. Pregnant women were excluded from the analysis and the sample included respondents above the age of 20 years who were medically examined and had laboratory tests. The analytical sample included 4693 US-born Blacks and 2968 foreign-born Blacks. We analyzed data using SAS 9.3 [26]. We weighted descriptive statistics and logistic regressions using the mobile examination centers survey weight provided by NHANES and utilized survey commands to account for the complex sampling design of the data.

**Dependent Variables** Systolic blood pressure, diastolic blood pressure, and 60-s pulse were measured by trained medical personnel. DBP and SBP were calculated by taking the average of two readings. We dichotomized each biomarker, using the 75th percentile to categorize respondents at greater risk of developing disease relative to the rest of the sample. Respondents with a reading at or above the 75th percentile were considered “high” risk (score = 1), with measures below the 75th percentile considered “low” risk (score = 0) for each biomarker. Individuals who reported taking medication for hypertension were coded in the high risk category when scoring blood pressure [27, 28].

**Measures**

Nativity was measured as a dichotomous variable indicating whether an individual was born outside or in the

USA (1 = foreign-born; 0 = US-born). NHANES defines length of time in the USA in years as a variable with 9 categories, which we recoded into three categories (1 = < 5 years; 2 = 5–19 years, 3 = > 20 years), consistent with other studies on immigrant health. Age was measured as a continuous variable defined as participants’ age in years at the time of the screening interview. Gender was based on self-reported information on whether a respondent was male or female (1 = woman; 0 = man). We defined education as a 4-level categorical variable (1 = < high school; 2 = high school diploma or general equivalency diploma; 3 = > high school diploma, some college, or an associate’s degree; 4 = > college degree). NHANES defines the poverty income ratio (PIR) as a ratio of family income to the federal poverty level based on family size in a calendar year with values ranging from 0 to 5. A PIR value of 1 is 100% of the federal poverty level, a

**Table 1** Weighted descriptive statistics for US-born Blacks and foreign-born Blacks by nativity: National Health and Nutrition Examination Survey, United States, 2001–2016

Characteristic	All (n <sup>a</sup> =7661) %	Foreign-born (n <sup>a</sup> =2968) %	US-born (n <sup>a</sup> =4693) %	P
Systolic blood pressure, mean (SD) <sup>b</sup>	126.06 (0.32)	126.41 (0.53)	125.92 (0.42)	0.816
Diastolic blood pressure, mean (SD) <sup>b</sup>	72.18 (0.26)	71.27 (0.42)	72.50 (0.29)	0.011
60-s pulse, mean (SD)	72.48 (0.21)	72.11 (0.37)	72.78 (0.24)	0.065
Age, mean (SD)	44.1 (0.29)	44.2 (0.53)	44.0 (0.34)	0.778
Gender				
Male	51.48	53.47	49.85	0.001
Female	48.52	46.53	50.15	
Length of US residence (years) <sup>c</sup>				
< 5	NA	24.14	NA	
5–19	NA	37.77	NA	
20 +	NA	38.09	NA	
Cigarette smoking status				
Current non-smoker	30.78	69.55	68.24	0.380
Current smoker	69.22	30.45	31.76	
Marital status				
Unmarried	33.07	37.29	30.77	< .0001
Married/living with partner	44.04	41.45	45.35	
Divorced/separated/widowed	22.89	21.26	23.88	
Poverty income ratio				
< 300% Federal poverty level	63.39	70.57	67.42	0.200
≥ 300% Federal poverty level	31.61	29.43	32.58	
Education				
Less than high school/no diploma	33.77	20.08	27.45	< .0001
High school diploma	19.07	26.57	26.08	
Some college or AA	24.71	35.33	32	
College graduate or above	22.45	18.02	14.47	

NA = Not Applicable

<sup>a</sup> Refers to unweighted totals

<sup>b</sup> Normal systolic blood pressure is < 120 and normal diastolic blood pressure is < 80, per American Heart Association Guidelines

<sup>c</sup> Length of US residence only applies to foreign-born respondents

**Table 2** Logistic regression results for diastolic blood pressure among Blacks by nativity status (US- or foreign-born): National Health and Nutrition Examination Survey, United States, 2001–2016

	Model 1 OR (95% CI)		Model 2 OR (95% CI)	
	All	Foreign	All	Foreign
Nativity				
Time in the USA				
20 + years	0.86 (0.75, 0.97)*	1.0 (ref)	1.52 (1.01, 2.30)*	1.0 (ref)
5–19 years		0.38 (0.20, 0.74)*		0.57 (0.28, 1.18)
<5 years		0.21 (0.09, 0.50)*		0.35 (0.14, 0.85)*
Age			1.07 (1.05, 1.08)*	1.05 (1.02, 1.08)*
Gender				
Female			1.0 (ref)	1.0 (ref)
Male			1.12 (0.74, 1.69)*	1.13 (0.61, 2.09)
Marital status				
Married/partnered			1.0 (ref)	1.0 (ref)
Divorced/widowed			0.84 (0.51, 1.37)	0.76 (0.41, 1.41)
Unmarried			0.71 (0.42, 1.20)	0.70 (0.34, 1.48)
Poverty income ratio				
> = 300% FPL				1.0 (ref)
< 300%FPL				0.93 (0.43, 2.02)
Education				0.61 (0.28, 1.35)
College grad or above				
Some college or AA				
High school grad				
No HS diploma/GED				
Smoking				
Non-smoker				
Smoker				
Model 3 OR (95% CI)				
	All	Foreign	US	US
Nativity				
Time in the USA				
20 + years	0.97 (0.85, 1.10)	1.0 (ref)		
5–19 years				
<5 years				
Age	1.07 (1.06, 1.08)*	1.05 (1.02, 1.09)*	1.06 (1.05, 1.08)*	1.07 (1.05, 1.09)*
Gender				
Female	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Male	1.06 (0.94, 1.19)	1.11 (0.54, 2.29)	1.29 (0.73, 2.28)	1.49 (0.86, 2.59)
Model 4 OR (95% CI)				
	All	Foreign	US	US
Nativity				
Time in the USA				
20 + years				
5–19 years				
<5 years				
Age	1.82 (1.15, 2.88)*	1.0 (ref)	1.06 (1.02, 1.10)*	1.07 (1.06, 1.09)*
Gender				
Female	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Male	1.21 (0.77, 1.91)	0.95 (0.40, 2.24)	1.21 (0.77, 1.91)	1.49 (0.86, 2.59)

**Table 2** (continued)

Marital status	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Married/partnered	0.94 (0.81, 1.10)	0.75 (0.33, 1.74)	1.02 (0.46, 2.26)	0.86 (0.45, 1.65)	0.64 (0.25, 1.60)	1.06 (0.41, 2.75)	
Divorced/widowed	0.79 (0.68, 0.90)*	0.84 (0.36, 1.93)	0.69 (0.31, 1.55)	0.81 (0.43, 1.53)	0.81 (0.30, 2.20)	0.69 (0.29, 1.67)	
Unmarried	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	
Poverty income ratio	1.04 (0.89, 1.21)	0.61 (0.25, 1.51)	0.74 (0.36, 1.49)	0.66 (0.38, 1.14)	0.55 (0.19, 1.62)	0.78 (0.41, 1.49)	
>= 300% FPL	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	
< 300%FPL	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	
Education	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	
College grad or above	1.15 (0.94, 1.40)	0.90 (0.37, 2.19)	1.21 (0.53, 2.74)	1.04 (0.58, 1.88)	0.85 (0.35, 2.03)	1.27 (0.54, 3.01)	
Some college or AA	1.19 (0.99, 1.44)	1.64 (0.71, 3.77)	0.72 (0.35, 1.48)	0.78 (0.45, 1.35)	1.11 (0.51, 2.41)	0.62 (0.33, 1.17)	
High school grad	1.12 (0.90, 1.40)	1.22 (0.54, 2.76)	1.52 (0.66, 3.46)	1.25 (0.64, 2.42)	1.09 (0.39, 3.05)	1.31 (0.53, 3.23)	
No HS diploma/GED							
Smoking							
Non-smoker							1.0 (ref)
Smoker							2.83 (0.70, 11.47)

OR = odds ratio, CI = confidence interval. \*P < .05

value of 2 is 200% of the federal poverty level, and so forth. We coded values of 4 and above as 5 to protect individuals' anonymity. We recoded this variable into 2 categories: between 0.00 and 2.99, and 3.00 and greater to reflect individuals who were at or below \$59,800 and those who were at or above \$59,801. Marital status was coded into a 3-level categorical variable (1 = never married; 2 = married or living with a partner; 3 = widowed, divorced, or separated). Smoking status was defined using questions asking about the average number of cigarettes smoked per day in the past 30 days. Current non-smokers were defined as those who had never smoked at least 100 cigarettes in their lifetime and did not smoke at the time of the interview. We designated individuals who smoked at least 100 cigarettes in their lifetime and currently smoked as current smokers (1 = smoker; 0 = non-smoker).

### Data Analysis

First, we compared the distribution of each biomarker between foreign- and the US-born Blacks utilizing *t* tests for continuous variables and Rao-Scott modified chi-square tests to account for the complex survey design for categorical variables. We then fit multivariable-adjusted logistic regression models to assess the association between nativity and each cardiovascular biomarker, adding covariates in a step-wise fashion across four models: Model 1, unadjusted; Model 2, adjusted for time in the USA (foreign-born only), age, gender, and marital status; Model 3, added PIR and education; and Model 4, fully adjusted (added smoking status). Because there are few variables that could affect nativity, and thus act as true confounders, adding potential mediating variables, such as PIR, education, and smoking, in a step-wise fashion, allowed use to evaluate changes in effect sizes with the addition of each variable. We tested models for the full sample as well as separate models for foreign- and the US-born Blacks to examine associations between sociodemographic variables and cardiovascular biomarkers in each population of interest.

### Results

Table 1 shows the stratified descriptive statistics for the sociodemographic variables and dependent variables by nativity. Both foreign- and the US-born had the same mean age of 44 years and had similar profiles with regard to the proportion of smokers and non-smokers. Compared with the US-born, a larger proportion of the foreign-born were male (54% vs 50%), unmarried (37% vs 31%), and had a college degree or above (18% vs 14%). A larger proportion of the US-born were female (50% vs 46%). There were no statistically significant differences in mean SBP, or 60-s pulse; however,

**Table 3** Logistic regression results for 60 second pulse among Blacks by Nativity status (US- or Foreign-Born): National Health and Nutrition Examination Survey, United States, 2001–2016

	Model 1 OR (95% CI)		Model 2 OR (95% CI)	
	Foreign	US	All	Foreign
Nativity				
Time in the USA			1.52 (0.91, 2.54)	
20 + years	1.0 (ref)			1.0 (ref)
5–19 years	0.96 (0.51, 1.79)			0.76 (0.38, 1.53)
< 5 years	1.19 (0.51, 2.77)			1.02 (0.40, 2.61)
Age			0.99 (0.98, 1.01)	0.98 (0.96, 1.00)
Gender				
Female			1.0 (ref)	1.0 (ref)
Male			0.44 (0.30, 0.65)*	0.32 (0.17, 0.60)*
Marital status				
Married/partnered			1.0 (ref)	1.0 (ref)
Divorced/widowed			1.01 (0.57, 1.79)	0.98 (0.47, 2.02)
Unmarried			1.46 (0.80, 2.65)	1.59 (0.76, 3.32)
Poverty income ratio				
> = 300% FPL				1.0 (ref)
< 300% FPL				1.09 (0.45, 2.66)
Education				1.29 (0.49, 3.43)
College grad or above				
Some college or AA				
High school grad				
No HS diploma/GED				
Smoking				
Non-smoker				
Smoker				

  

	Model 3 OR (95% CI)		Model 4 OR (95% CI)	
	Foreign	US	All	Foreign
Nativity				
Time in the USA			1.78 (1.02, 3.10)**	
20 + years	1.0 (ref)			1.0 (ref)
5–19 years	0.77 (0.33, 1.82)			0.81 (0.33, 1.99)
< 5 years	0.71 (0.29, 1.77)			0.95 (0.39, 2.32)
Age			0.99 (0.97, 1.01)	0.98 (0.95, 1.01)
Gender				
Female			1.0 (ref)	1.0 (ref)
Male			0.49 (0.31, 0.79)*	1.95 (0.85, 4.50)



**Table 3** (continued)

Marital status	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Married/partnered	1.02 (0.58, 1.81)	1.05 (0.52, 2.11)	0.98 (0.39, 2.47)	1.23 (0.66, 2.32)	1.22 (0.56, 2.67)	1.20 (0.42, 3.42)	1.20 (0.42, 3.42)	1.20 (0.42, 3.42)	1.20 (0.42, 3.42)	1.20 (0.42, 3.42)
Divorced/widowed	1.77 (0.93, 3.40)	2.20 (0.97, 5.02)	1.34 (0.50, 3.62)	1.87 (0.94, 3.71)	2.20 (0.91, 5.32)	1.48 (0.52, 4.22)	1.48 (0.52, 4.22)	1.48 (0.52, 4.22)	1.48 (0.52, 4.22)	1.48 (0.52, 4.22)
Unmarried	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Poverty income ratio	1.28 (0.75, 2.17)	2.04 (0.89, 4.75)	0.91 (0.47, 1.77)	1.08 (0.59, 1.95)	1.95 (0.85, 4.50)	0.61 (0.27, 1.41)	0.61 (0.27, 1.41)	0.61 (0.27, 1.41)	0.61 (0.27, 1.41)	0.61 (0.27, 1.41)
> = 300% FPL	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
< 300%FPL	1.28 (0.75, 2.17)	2.04 (0.89, 4.75)	0.91 (0.47, 1.77)	1.08 (0.59, 1.95)	1.95 (0.85, 4.50)	0.61 (0.27, 1.41)	0.61 (0.27, 1.41)	0.61 (0.27, 1.41)	0.61 (0.27, 1.41)	0.61 (0.27, 1.41)
Education	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
College grad or above	1.28 (0.68, 2.43)	0.90 (0.38, 2.16)	1.62 (0.65, 4.08)	1.29 (0.67, 2.49)	0.82 (0.33, 2.02)	1.92 (0.74, 4.99)	1.92 (0.74, 4.99)	1.92 (0.74, 4.99)	1.92 (0.74, 4.99)	1.92 (0.74, 4.99)
Some college or AA	1.68 (0.74, 3.79)	1.35 (0.44, 4.08)	1.90 (0.57, 6.33)	1.40 (0.62, 3.16)	1.13 (0.45, 2.85)	1.32 (0.37, 4.76)	1.32 (0.37, 4.76)	1.32 (0.37, 4.76)	1.32 (0.37, 4.76)	1.32 (0.37, 4.76)
High school grad	1.03 (0.46, 2.31)	0.58 (0.16, 2.14)	1.75 (0.66, 4.64)	1.02 (0.43, 2.46)	0.53 (0.14, 1.99)	2.07 (0.61, 7.01)	2.07 (0.61, 7.01)	2.07 (0.61, 7.01)	2.07 (0.61, 7.01)	2.07 (0.61, 7.01)
No HS diploma/GED	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Smoking	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Non-smoker	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Smoker	0.49 (0.16, 1.54)	0.54 (0.10, 2.96)	0.54 (0.10, 2.96)	0.54 (0.10, 2.96)	0.54 (0.10, 2.96)	0.54 (0.10, 2.96)	0.54 (0.10, 2.96)	0.54 (0.10, 2.96)	0.54 (0.10, 2.96)	0.54 (0.10, 2.96)

OR = odds ratio, CI = confidence interval. \*P < .05

the US-born had a statistically significantly higher mean DBP compared with the foreign-born.

### Full Sample

Tables 2, 3, and 4 show the results of logistic regression for mean SBP, mean DBP, and 60-s pulse. For the full sample, foreign-born Blacks were more likely to have high mean DBP and 60-s pulse compared with the US-born individuals. In the final model, foreign-born Blacks were almost twice as likely to have a high mean DBP compared with the US-born individuals (odds ratio [OR]= 1.82; 95% confidence interval [CI]= 1.15, 2.88). Foreign-born Blacks were more likely to have a high mean 60-s pulse, (OR = 1.78; 95% CI = 1.02, 3.10).

Of note, among the sociodemographic variables, compared with younger individuals, older individuals were more likely to have high DBP (OR = 1.07; 95% CI = 1.05, 1.09) and high mean SBP (OR = 1.10; 95% CI = 1.08, 1.13). Men were 58% less likely to have a risk of high 60-s pulse compared with women (OR = 0.42; 95% CI = 0.27, 0.66).

### The US-Born

Older individuals were more likely to have high mean SBP and DBP compared with younger individuals (OR = 1.11; 95% CI = 1.07, 1.14), (OR = 1.07; 95% CI = 1.06, 1.09) respectively. Men were less than half as likely to have a high 60-s pulse compared with women (OR = 0.576; 95% CI = 0.496, 0.670). There were no other significant effects observed between sociodemographic variables or any outcome for the US-born Blacks.

### Foreign-Born

Foreign-born Blacks living in the USA for less than 5 years were 62% less likely to have high SBP compared with those in the USA for 20 years or more (OR = 0.38; 95% CI = 0.18, 0.80). However, this effect did not remain significant when the other sociodemographic variables were added to the model. An effect for age was observed in the final mean SBP and mean DBP models when all variables were added to the model, such that older individuals were more likely to have high mean SBP and DBP compared with younger individuals (OR = 1.11; CI = 1.07, 1.15; OR = 1.06; CI = 1.02, 1.10). Foreign-born men were less likely to have a high 60-s pulse compared with women (OR = 0.37; CI = 0.20, 0.70). There were no other significant effects observed.

### Discussion

This study aimed to examine nativity-based differences in the cardiovascular biomarkers mean SBP, mean DBP, and 60-s

**Table 4** Logistic regression results for systolic blood pressure among Blacks by nativity status (US- or foreign-born): National Health and Nutrition Examination Survey, United States, 2001–2016

Nativity	Model 1 OR (95% CI)		Model 2 OR (95% CI)	
	All	US	All	Foreign
	0.89 (0.61, 1.30)		1.29 (0.83, 2.00)	
Time in the USA				
20 + years		1.0 (ref)		1.0 (ref)
5–19 years		0.41 (0.21, 0.81)*		0.87 (0.43, 1.77)
< 5 years		0.38 (0.18, 0.80)*		1.10 (0.49, 2.47)
Age			1.10 (1.08, 1.12)*	1.10 (1.06, 1.13)*
Gender				
Female			1.0 (ref)	1.0 (ref)
Male			1.08 (0.71, 1.65)	0.80 (0.43, 1.51)
Marital status				
Married/partnered			1.0 (ref)	1.0 (ref)
Divorced/widowed			0.80 (0.46, 1.39)	0.80 (0.35, 1.81)
Unmarried			0.99 (0.50, 1.96)	0.70 (0.30, 1.64)
Poverty income ratio				
> = 300% FPL				1.0 (ref)
< 300% FPL				0.78 (0.36, 1.69)
Education				1.41 (0.51, 3.86)
College grad or above				
Some college or AA				
High school grad				
No HS diploma/GED				
Smoking				
Non-smoker				
Smoker				
			Model 4 OR (95% CI)	
Nativity	Model 3 OR (95% CI)		All	Foreign
	All	US	1.41 (0.86, 2.31)	
	1.31 (0.83, 2.06)			
Time in the USA				
20 + years				1.0 (ref)
5–19 years				1.38 (0.55, 3.45)
< 5 years				1.66 (0.55, 5.08)
Age			1.10 (1.08, 1.12)*	1.11 (1.07, 1.15)*
Gender				
Female			1.0 (ref)	1.0 (ref)
Male			1.31 (0.83, 2.05)	0.75 (0.33, 1.70)
			1.10 (1.09, 1.10)*	1.11 (1.07, 1.14)*
			1.0 (ref)	1.0 (ref)
			1.23 (0.74, 2.04)	1.60 (0.86, 2.95)

**Table 4** (continued)

Marital status	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Married/partnered	0.89 (0.48, 1.66)	0.86 (0.32, 2.33)	1.08 (0.88, 1.31)	0.87 (0.43, 1.77)	0.77 (0.24, 2.48)	0.82 (0.32, 2.09)	0.82 (0.32, 2.09)
Divorced/widowed	1.22 (0.61, 2.42)	0.92 (0.38, 2.22)	0.88 (0.70, 1.10)	1.16 (0.55, 2.44)	0.95 (0.36, 2.48)	1.55 (0.50, 4.77)	1.55 (0.50, 4.77)
Unmarried	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Poverty income ratio	0.72 (0.39, 1.32)	0.60 (0.25, 1.42)	1.19 (0.99, 1.44)	0.71 (0.37, 1.36)	0.51 (0.21, 1.22)	0.83 (0.32, 2.16)	0.83 (0.32, 2.16)
> = 300% FPL	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
< 300%FPL	1.70 (0.86, 3.38)	1.78 (0.70, 4.52)	1.14 (0.88, 1.48)	1.67 (0.83, 3.33)	1.57 (0.56, 4.40)	1.66 (0.60, 4.57)	1.66 (0.60, 4.57)
Education	1.53 (0.73, 3.23)	1.80 (0.58, 5.67)	1.20 (0.92, 1.56)	1.07 (0.52, 2.21)	0.90 (0.24, 3.37)	1.12 (0.46, 2.75)	1.12 (0.46, 2.75)
College grad or above	1.31 (0.68, 2.52)	1.02 (0.39, 2.65)	1.10 (0.83, 1.45)	1.16 (0.57, 2.38)	0.79 (0.29, 2.17)	1.52 (0.59, 3.86)	1.52 (0.59, 3.86)
Some college or AA	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
High school grad	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
No HS diploma/GED	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Smoking	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Non-smoker	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Smoker	1.41 (0.86, 2.31)	1.41 (0.86, 2.31)	1.41 (0.86, 2.31)	1.41 (0.86, 2.31)	2.73 (0.77, 9.70)	0.92 (0.31, 2.75)	0.92 (0.31, 2.75)

OR = odds ratio, CI = confidence interval. \*P < .05

pulse. We also aimed to describe differences in associations between the sociodemographic variables under study and these outcomes among foreign- vs the US-born samples. Our findings suggest that there were no nativity-based differences in mean SBP; however, compared with the US-born, the foreign-born were more likely to have higher risks of mean DBP and 60-s pulse. This finding contradicts past views on the HIE and the foreign-born health advantage but is in line with more recent re-examinations of the HIE, highlighting that Black immigrants have higher rates of hypertension than their US-born counterparts [6, 19–21]. More current views of Black migration points to the notion that Africa and other Black-majority nations are becoming more urbanized with an increase in processed food consumption and less exercise. As a result, we can speculate that recent cohorts of immigrants may not be arriving in the USA as healthy as previous cohorts.

Due to the high prevalence of hypertension in the USA and globally and its associated morbidities as age increases, hypertension has become the dominant risk factor for cardiovascular disease in older individuals [29]. We found significant age differences in SBP and DBP and gender differences in 60-s pulse across the full sample and in the separate foreign- and the US-born models. We also consistently observed an increase in the risk of mean SBP and mean DBP with increased age in the full sample and among the foreign- and the US-born in separate models. These findings have been confirmed in other, larger epidemiological studies although there is a need for studies focused on immigrants from African and other Black-majority nations [24, 30].

Although time spent in the USA was a predictor of mean SBP among the foreign-born sample, this significant result disappeared in the final model. Due to the cross-sectional nature of the data, we cannot determine how covariates may have changed post-immigration. For example, changes in an individual's income may have mediated the association between time in the USA and SBP, which would attenuate the effect observed in unadjusted models. We found that individuals living in the USA for less than 5 years were less likely to have high SBP compared with those in the USA for 20 years or more. This is consistent with other studies demonstrating that over time, the foreign-born lose any health advantage that they may have possessed upon arrival as they adjust to the sociocultural environment of the USA. Furthermore, studies of other immigrant groups in industrialized societies have shown that each additional year of US residence is associated with increased odds of hypertension [30–32].

### Limitations

This study is not without limitations. First, the study was cross-sectional in nature and therefore limits our ability to make conclusive statements regarding causality. While we cannot rule out pre-existing conditions, because our outcomes

were objective, we can be assured that immigration occurred prior to the collection of cardiovascular outcomes. Second, we were unable to examine birth country specific data which would be useful to further disaggregate the effects of nativity. Although hypertension and cardiovascular disease are a global public health concern, some nations are experiencing higher incidences than others, which may impact our assessment of nativity-based differences in the cardiovascular risk biomarkers. Lastly, our sociodemographic and nativity information was self-reported, which is vulnerable to recall and information bias.

### Public Health Implications

Our results suggest that there is no foreign-born health advantage for any of the separate cardiovascular risk biomarkers evaluated in this study. Moreover, foreign-born Blacks were at higher risk of elevated mean DBP and 60-s pulse than their US-born counterparts. Among both populations, advanced age seems to work similarly and increases the risk of having higher mean SBP and mean DBP. Importantly, our results highlight that there is no nativity-based difference in mean SBP, the more commonly used measure in evaluating and controlling hypertension.

We considered that although hypertension is a useful measure of cardiovascular function, examining the biomarkers separately provides beneficial information about how various sociodemographic variables work among the foreign- and the US-born. Researchers should consider disaggregating hypertension data into its individual components in order to fully capture the mechanisms at play. Future research should continue to examine the HIE in larger Black immigrant populations, as well as the utility of the aggregate hypertension measure as a clinical marker for cardiovascular risk, as opposed to its individual components [33].

### Compliance with Ethical Standards

**Conflict of Interest** The authors declare that they have no conflict of interest.

**Ethical Approval** This article does not include any studies with human participants or animals performed by any of the authors.

**Informed Consent** No protocol approval for informed consent was required, as the data are secondary data and publicly available.

### References

1. U.S. Census Bureau. American community survey. American FactFinder; 2015.
2. Department of Homeland Security (DHS), Office of Immigration Statistics. 2015 yearbook of immigration statistics. Washington, DC: DHS Office of Immigration Statistics; 2016.
3. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Borden WB, et al. Heart disease and stroke statistics—2013 update: a report from the American Heart Association. *Circulation*. 2013;127(1):e6–e245. <https://doi.org/10.1161/CIR.0b013e31828124ad>.
4. Mensah GA, Mokdad AH, Ford ES, Greenlund KJ, Croft JB. State of disparities in cardiovascular health in the United States. *Circulation*. 2005;111:1233–41.
5. Wang X, Poole JC, Treiber FA, Harshfield GA, Hanevold CD, Snieder H. Ethnic and gender differences in ambulatory blood pressure trajectories: results from a 15-year longitudinal study in youth and young adults. *Circulation*. 2006;114:2780–7.
6. Commodore-Mensah Y, Hill M, Allen J, Cooper LA, Blumenthal R, Agyemang C, et al. Sex differences in cardiovascular disease risk of Ghanaian- and Nigerian-born west African immigrants in the United States: the afro-cardiac study. *J Am Heart Assoc*. 2016. <https://doi.org/10.1161/JAHA.115.002385>.
7. Levitt NS, Steyn K, Dave J, Bradshaw D. Chronic noncommunicable diseases and HIV-AIDS on a collision course: relevance for health care delivery, particularly in low-resource settings—insights from South Africa. *Am J Clin Nutr*. 2011;94:1690S–6S.
8. Levitt NS. Diabetes in Africa: epidemiology, management and healthcare challenges. *Heart*. 2008;94:1376–82.
9. International Diabetes Federation. IDF Diabetes Atlas, 6th edn. 2013. Available at [www.idf.org/diabetesatlas](http://www.idf.org/diabetesatlas).
10. Moran A, Forouzanfar M, Sampson U, Chugh S, Feigin V, Mensah G. The epidemiology of cardiovascular diseases in sub-Saharan Africa: the global burden of diseases, injuries and risk factors 2010 study. *Prog Cardiovasc Dis*. 2013;5:234–9.
11. Muthuri SK, Oti SO, Lilford RJ, Oyebo O. Salt reduction interventions in sub-Saharan Africa: a systematic review. *PLoS One*. 2016;11:e0149680.
12. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *Lancet*. 2005;365:217–23.
13. Vidal-Petiot E, Ford I, Greenlaw N, Ferrari R, Fox KM, Tardif JC, et al. Investigators. Cardiovascular event rates and mortality according to achieved systolic and diastolic blood pressure in patients with stable coronary artery disease: an international cohort study. *Lancet*. 2016;29(388):2142–52.
14. van den Hoogen PC, Feskens EJ, Nagelkerke NJ, Menotti A, Nissinen A, Kromhout D. The relation between blood pressure and mortality due to coronary heart disease among men in different parts of the world. Seven Countries Study Research Group. *N Engl J Med*. 2000. <https://doi.org/10.1056/NEJM200001063420101>.
15. Rothwell PM, Howard SC, Dolan E, O'Brien E, Dobson JE, Dahlöf B, et al. Prognostic significance of visit-to-visit variability, maximum systolic blood pressure, and episodic hypertension. *Lancet*. 2010;375:895–905.
16. Franklin SS, Larson MG, Khan SA, Wong ND, Leip EP, Kannel WB, et al. Does the relation of blood pressure to coronary heart disease risk change with aging? The Framingham Heart Study. *Circulation*. 2001;103(9):1245–9.
17. Brown AG, Houser RF, Mattei J, Mozaffarian D, Lichtenstein AH, Foltz SC. Hypertension among US-born and foreign-born non-Hispanic blacks: NHANES 2003–2014 data. *J Hypertens*. 2017;35:2380–7.
18. Wilson TW, Hollifield LR, Grim CE. Systolic blood pressure levels in black populations in sub-Saharan Africa, the West Indies, and the United States: a meta-analysis. *Hypertension*. 1991;18:187–91.
19. Ukegbu UJ, Castillo DC, Knight MG, Ricks M, Miller BV, Onumah BM, et al. Metabolic syndrome does not detect metabolic risk in African men living in the US. *Diabetes Care*. 2011;34:2297–9.
20. Yu SS, Ramsey NL, Castillo DC, Ricks M, Sumner AE. Triglyceride-based screening tests fail to recognize cardiometabolic

- disease in African immigrant and African-American men. *Metab Syndr Relat Disord*. 2013;11:15–20.
21. O'Connor MY, Thoreson CK, Ricks M, Courville AB, Thomas F, Yao J, et al. Worse cardiometabolic health in African immigrant men than African American men: reconsideration of the healthy immigrant effect. *Metab Syndr Relat Disord*. 2014;12:347–53.
  22. Commodore-Mensah Y, Himmelfarb CD, Agyemang C, Sumner AE. Cardiometabolic health in African immigrants to the United States: a call to re-examine research on African-descent populations. *Ethn Dis*. 2015;25:373–80.
  23. Bidulescu A, Francis DK, Ferguson TS, Bennett NR, Hennis AJ, Wilks R, et al. Disparities in hypertension among black Caribbean populations: a scoping review by the US Caribbean Alliance for Health Disparities Research Group (USCAHDR). *Int J Equity Health*. 2015;14:125.
  24. Vokonas PS, Kannel WB, Cupples LA. Epidemiology and risk of hypertension in the elderly: the Framingham study. *J Hypertens Suppl*. 1988;6(1):S3–9.
  25. Centers for Disease Control and Prevention. National Health and Nutrition Examination Survey Data. 2012. Available at: [www.cdc.gov/nchs/nhanes/nhanes\\_questionnaires.htm](http://www.cdc.gov/nchs/nhanes/nhanes_questionnaires.htm). Accessed 20 June 2018.
  26. SAS/STAT. Version 9.3. [computer program]. Cary: SAS Institute; 2008.
  27. Geronimus AT, Hicken M, Keene D, Bound J. Weathering and age patterns of allostatic load scores among blacks and whites in the United States. *Am J Public Health*. 2006;96:826–33.
  28. Crimmins EM, Kim JK, Alley DE, Karlamangla A, Seeman T. Hispanic paradox in biological risk profiles. *Am J Public Health*. 2007;97:1305–10.
  29. Zhou B, Bentham J, Di Cesare M, Bixby H, Danaei G, Cowan MJ, et al. Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19·1 million participants. *Lancet*. 2017;389:37–55.
  30. Lloyd-Jones DM, Evans JC, Levy D. Hypertension in adults across the age spectrum: current outcomes and control in the community. *JAMA*. 2005;294:466–72.
  31. Moran A, Roux AV, Jackson SA, Kramer H, Manolio TA, Shrager S, et al. Acculturation is associated with hypertension in a multiethnic sample. *Am J Hypertens*. 2007;20:354–63.
  32. Steffen PR, Smith TB, Larson M, Butler L. Acculturation to Western society as a risk factor for high blood pressure: a meta-analytic review. *Psychosom Med*. 2006;68:386–97.
  33. Flack JM, Ferdinand KC, Nasser SA. Epidemiology of hypertension and cardiovascular disease in African Americans. *J Clin Hypertens*. 2003;5(1):5–11.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.